Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Then $\frac{ds_1}{d\theta}=a(1-e^2\cos^2\theta)^{\frac{1}{2}}$ and $\frac{ds_2}{d\phi}=a(1-e^2\cos^2\phi)^{\frac{1}{2}}$.

Then the required average is

$$A=\frac{H}{S^2}\int_0^{\frac{1}{2}\pi}\int_0^{2\pi}Kds_1ds_2=\frac{4a^2}{S^2}\int_0^{\frac{1}{2}\pi}\int_0^{2\pi}[a^2(\cos\phi-\cos\theta)^2+\frac{b^2}{a^2}(\sin\phi-\sin\theta)^2]^{\frac{1}{2}}\times$$

$$(1-e^2\cos^2\theta)^{\frac{1}{2}}(1-e^2\cos^2\phi)^{\frac{1}{2}}\,d\theta d\phi.$$

This equation cannot be integrated in general terms.

Solved in the same manner by *G. B. M. ZERR*,

41. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Irving College, Mechanicsburg, Pennsylvania.

A line is drawn at random across the chord and given arc of a circular segment. Find the mean area of the divisions.

Solution by G. B. M. ZERR, A. M., Ph. D., Texarkana, Arkansas-Texas.

Let A=area of given segment, A_1, A_2 mean areas of the two divisions.
$\therefore A_1+A_2=A$.
But, since the line is a random line, $A_1=A_2$.
$\therefore A_1=A_2=\frac{1}{2}A$.

Also solved by *HENRY HEATON*.

42. Proposed by CHARLES E. MYERS, Canton, Ohio.

A attends church 4 Sundays out of 5; B, 5 Sundays out of 6; and C, 6 Sundays out of 7. What is the probability of an event that A and B will be at church and C will not?

Solution by G. B. M. ZERR, A. M., Ph. D., Texarkana, Arkansas-Texas, and B. F. FINKEL, A. M., Professor of Mathematics and Physics, Drury College, Springfield, Missouri.

The chance that A attends church$=\frac{4}{5}$.
The chance that B attends church$=\frac{5}{6}$.
The chance that C attends church$=\frac{6}{7}$.
The chance that A is not at church$=\frac{1}{5}$.
The chance that B is not at church$=\frac{1}{6}$.
The chance that C is not at church$=\frac{1}{7}$.
The chance that A and B attend and C not$=p_1=\frac{4}{5}.\frac{5}{6}.\frac{1}{7}=\frac{2}{21}$.
The chance that A and C attend and B not$=p_2=\frac{4}{5}.\frac{6}{7}.\frac{1}{6}=\frac{4}{35}$.
The chance that B and C attend and A not$=p_3=\frac{5}{6}.\frac{6}{7}.\frac{1}{5}=\frac{1}{7}$.
The chance that A attends and B and C not$=p_4=\frac{4}{5}.\frac{1}{6}.\frac{1}{7}=\frac{4}{210}$.
The chance that B attends and A and C not$=p_5=\frac{1}{5}.\frac{5}{6}.\frac{1}{7}=\frac{1}{42}$.
The chance that C attends and A and B not$=p_6=\frac{1}{5}.\frac{1}{6}.\frac{6}{7}=\frac{1}{35}$.
The chance that A, B and C attend$=p_7=\frac{4}{5}.\frac{5}{6}.\frac{6}{7}=\frac{4}{7}$.
The chance that A, B and C do not attend$=p_8=\frac{1}{5}.\frac{1}{6}.\frac{1}{7}=\frac{1}{210}$.
p_1=probability required.
Also $p_1+p_2+p_3+p_4+p_5+p_6+p_7+p_8=1$.

Also solved by *HENRY HEATON*.